

Effect of Availability Bias and Reflective Reasoning on Diagnostic Accuracy Among Internal Medicine Residents

Sílvia Mamede, MD, PhD

Tamara van Gog, MSc, PhD

Kees van den Berge, MD

Remy M. J. P. Rikers, Msc, PhD

Jan L. C. M. van Saase, MD, PhD

Coen van Guldener, MD, PhD

Henk G. Schmidt, MSc, PhD

A MAJOR AIM OF EVERY CLINICAL teacher is to foster the quality of students' and residents' clinical reasoning, one of the most important factors affecting individual physicians' performance.¹ Diagnostic errors constitute a substantial portion of preventable medical mistakes,² and they have been attributed to a large extent to faulty clinical reasoning.¹ The development of educational strategies to minimize flaws in clinical reasoning depends on a better understanding of their underlying cognitive mechanisms.

Cognitive biases are a source of flaws in reasoning processes.³ At least 40 types of biases that may affect clinical reasoning have been described.^{4,5} A prime example is a biased use of the availability heuristic (the tendency to weigh likelihood of things by how easily they are recalled), which may erroneously lead a physician to consider a diagnosis more frequently and judge it as more likely if it comes to mind more easily.^{4,6} Relying on availability is often helpful during reasoning because things that come to mind easily generally do occur more frequently. However, a serious problem may arise when

For editorial comment see p 1233.

Context Diagnostic errors have been associated with bias in clinical reasoning. Empirical evidence on the cognitive mechanisms underlying biases and effectiveness of educational strategies to counteract them is lacking.

Objectives To investigate whether recent experience with clinical problems provokes availability bias (overestimation of the likelihood of a diagnosis based on the ease with which it comes to mind) resulting in diagnostic errors and whether reflection (structured reanalysis of the case findings) counteracts this bias.

Design, Setting, and Participants Experimental study conducted in 2009 at the Erasmus Medical Centre, Rotterdam, with 18 first-year and 18 second-year internal medicine residents. Participants first evaluated diagnoses of 6 clinical cases (phase 1). Subsequently, they diagnosed 8 different cases through nonanalytical reasoning, 4 of which had findings similar to previously evaluated cases but different diagnoses (phase 2). These 4 cases were subsequently diagnosed again through reflective reasoning (phase 3).

Main Outcome Measures Mean diagnostic accuracy scores (perfect score, 4.0) on cases solved with or without previous exposure to similar problems through nonanalytical (phase 2) or reflective (phase 3) reasoning and frequency that a potentially biased (ie, phase 1) diagnosis was given.

Results There were no main effects, but there was a significant interaction effect between "years of training" and "recent experiences with similar problems." Results consistent with an availability bias occurred for the second-year residents, who scored lower on the cases similar to those previously encountered (1.55; 95% confidence interval [CI], 1.15-1.96) than on the other cases (2.19; 95% CI, 1.73-2.66; $P=.03$). This pattern was not seen among the first-year residents (2.03; 95% CI, 1.55-2.51 vs 1.42; 95% CI, 0.92-1.92; $P=.046$). Second-year residents provided the phase 1 diagnosis more frequently for phase 2 cases they had previously encountered than for those they had not (mean frequency per resident, 1.44; 95% CI, 0.93-1.96 vs 0.72; 95% CI, 0.28-1.17; $P=.04$). A significant main effect of reasoning mode was found: reflection improved the diagnoses of the similar cases compared with nonanalytical reasoning for the second-year residents (2.03; 95% CI, 1.49-2.57) and the first-year residents (2.31; 95% CI, 1.89-2.73; $P=.006$).

Conclusion When faced with cases similar to previous ones and using nonanalytical reasoning, second-year residents made errors consistent with the availability bias. Subsequent application of diagnostic reflection tended to counter this bias; it improved diagnostic accuracy in both first- and second-year residents.

JAMA. 2010;304(11):1198-1203

www.jama.com

this first impression is wrong, because physicians often become anchored in their initial hypothesis, looking for confirming evidence to support their initial diagnosis, underestimating evidence against it, and therefore failing to adjust their initial impression in light of all available information.^{4,7}

Author Affiliations: Departments of Psychology (Drs Mamede, van Gog, Rikers, and Schmidt) and Internal Medicine, Erasmus Medical Centre (Drs van den Berge and van Saase), Erasmus University Rotterdam; and Department of Internal Medicine, Amphia Hospital, Breda (Dr van Guldener), the Netherlands. **Corresponding Author:** Sílvia Mamede, MD, MPH, PhD, Department of Psychology, Erasmus University Rotterdam, Burgemeester Oudlaan 50, Rotterdam, 3062 PA, the Netherlands (mamede@fsw.eur.nl).

The scientific literature on the availability bias in medicine is mainly descriptive. Some correlational studies⁸⁻¹¹ suggest that it occurs, but these do not allow causal inferences to be made. Experimental research is required to provide direct evidence for availability bias in medical diagnosis but, to the best of our knowledge, it is lacking. Moreover, if documented, it is perhaps even more important to medical education and practice to investigate ways in which availability bias can be counteracted.

Expertise might play a role in bias. Experienced physicians tend to rely more on nonanalytical (or System 1) reasoning based on pattern recognition to diagnose routine problems; this is a rapid, largely unconscious diagnostic approach. Although effective and highly efficient in most cases, it might be more easily affected by biases.^{12,13} One way to counteract biases, suggested by studies in psychology,^{4,14} is to induce physicians to adopt more reflective (or analytical, also referred to as System 2) reasoning, which comprises careful, effortful consideration of findings in a case, or to combine nonanalytical and analytical reasoning.¹⁵

Therefore, we investigated whether availability bias occurs when physicians diagnose cases that have clinical manifestations similar to those of recently encountered cases and if so, whether reflection could counteract this bias. Because nonanalytical reasoning develops in association with clinical experience, we also investigated whether there would be a difference in degree of bias between residents in the first and second year of a residency program. We hypothesized that (1) recent experiences with clinical problems would generate an availability bias when physicians nonanalytically diagnose subsequent cases of similar diseases; (2) more experienced residents would be more prone to this bias; and (3) reflective reasoning would counteract this bias and improve diagnostic accuracy.

METHODS

Overview

This experiment consisted of 3 phases conducted sequentially in a single ses-

Table 1. Clinical Cases Used in Each Phase of the Study

	Phase 1 Exposure (Diagnosis Evaluation Task)	Phase 2 Nonanalytical Diagnostic Reasoning	Phase 3 Reflective Diagnostic Reasoning
Set 1	Case A: acute viral hepatitis	Cases similar to case A ^a Liver cirrhosis Primary sclerosing cholangitis	Cases similar to case A Liver cirrhosis Primary sclerosing cholangitis
	Case B: inflammatory bowel disease	Cases similar to case B ^a Celiac disease Pseudomembranous colitis	Cases similar to case B Celiac disease Pseudomembranous colitis
	Neutral cases Meningitis Pyelonephritis Pneumonia Hyperthyroidism	Cases similar to case C ^b Acute viral pericarditis Aortic dissection	
Set 2	Case C: Acute myocardial infarction	Cases similar to case C ^a Acute viral pericarditis Aortic dissection	Cases similar to case C Acute viral pericarditis Aortic dissection
	Case D: Wernicke encephalopathy	Cases similar to case D ^a Neurosyphilis Vitamin B ₁₂ deficiency	Cases similar to case D Neurosyphilis Vitamin B ₁₂ deficiency
	Neutral cases Meningitis Pyelonephritis Pneumonia Hyperthyroidism	Cases similar to case A ^b Liver cirrhosis Primary sclerosing cholangitis	
		Cases similar to case B ^b Celiac disease Pseudomembranous colitis	

^aCases potentially subject to bias.

^bCases not subject to bias.

sion (TABLE 1). *Phase 1*, exposure, required participants to evaluate the accuracy of a diagnosis provided for 6 different cases. *Phase 2*, nonanalytical diagnosis, required participants to diagnose 8 new cases, 4 of which had clinical manifestations that were similar to 2 of the diseases encountered in phase 1. This was expected to induce an availability bias for those 4 cases and reduce diagnostic accuracy. *Phase 3*, reflective diagnosis, required participants to reflect on the diagnosis of the 4 cases that could have been influenced by an availability bias in phase 2. This was expected to overrule the bias and lead to more accurate diagnoses.

Participants

Thirty-six out of 42 eligible internal medicine residents (participation rate, 85.7%) from the Erasmus Medical Centre, Faculty of Medicine, Erasmus University Rotterdam (mean [SD] age, 29.5

[2.1] years) volunteered to participate in this study. Eighteen were in their first and 18 were in their second year of the residency program. The study took place during an educational meeting held in September 2009; the academic year starts in January for the majority of the residents. Participants did not receive any compensation or other incentives. The nonparticipants were either doing shifts or on holidays. The ethics review committee from the Department of Psychology, Erasmus University Rotterdam provided approval for this study. Because the nature of the study prevented prior disclosure of its objectives, oral consent was obtained after informing participants about their tasks. Debriefing was provided later.

Procedure

In total, 16 written clinical cases were used in this study (Table 1). Cases consisted of a brief description of a patient's medical history, signs and symp-

Box. Example of a Case (Diagnosis: Celiac Disease)

A 27-year-old woman presented with 11-month duration of complaints of diarrhea, flatulence, and episodes of abdominal cramps. She has had stools 5 to 6 times a day, and has often woken up during the night for defecation. The feces are voluminous and soft without mucus, blood, or pus. The abdominal cramps are more severe just before defecation, after which they become less painful. The patient is fatigued and has experienced a 5-kg weight loss over the past 11 months. She also noticed red spots on her skin. She says that she has not had fever or joint pains. The patient consulted a physician 4 months ago as well. The physician prescribed ferrous sulfate for anemia, which she has been using until now. Family history: her father was treated for lung tuberculosis 20 years ago.

Physical Examination

Young, somewhat emaciated woman of otherwise healthy appearance. BP: 110/70; pulse: 80/min; temperature: 36°C. Mucocutaneous paleness (+/4). No other abnormalities.

Laboratory Tests

Hemoglobin: 9 g/dL; hematocrit: 34%; Mean corpuscular volume: 74 fL; serum iron: 45 mg/dL (normal, 50-170 mg/dL); calcium: 8.1 mg/dL (normal, 8.6-10 mg/dL); albumin: 3.2 g/dL (normal, 3.4-4.8 g/dL); Alanine aminotransferase test: 38 U/L; Aspartate aminotransferase test: 25 U/L; Prothrombin time 24 seconds (normal, 12-22 seconds). Feces revealed no worm eggs, no parasites, no white cells; stool fat level was 12g /24 h (normal, <7g/24 h), D-xylose test was positive. Human immunodeficiency virus antibodies: negative. Tuberculosis skin test (PPD): 5 mm.

Imaging Tests

Chest x-ray: no abnormalities; colonoscopy: no abnormalities.

To convert alanine aminotransferase from U/L to $\mu\text{katal}(\text{kat})/\text{L}$, multiply by 0.0167; aspartate aminotransferase from U/L to μkat , multiply by 0.0167; iron from $\mu\text{g}/\text{dL}$ to $\mu\text{mol}/\text{L}$, multiply by 0.179.

toms, and tests results (example case shown in (BOX)). All cases were based on real patients with a confirmed diagnosis. They were prepared by experts in internal medicine and used in previous studies with internal medicine residents.^{16,17} The cases were presented to participants in a booklet (1 for each phase) in a random sequence.

In phase 1, each case had a diagnosis listed, and participants had to rate the likelihood (as percentage) that the indicated diagnosis was correct. The provided diagnosis was always correct, but participants were not aware of this, nor did they receive feedback on their likelihood ratings. This phase consisted of 6 cases: 4 neutral cases and 2 cases of diseases that have signs and symptoms also frequently encountered in 2 other diseases presented in phase 2 (Table 1). For example, a patient with cirrhosis or primary scleros-

ing cholangitis (phase 2) may present with signs and symptoms similar to acute viral hepatitis (phase 1). To minimize potential influence of case specificity or difficulty, we used 2 booklets with different sets of cases in phase 1; participants randomly received either set 1 or set 2. In each set, the similar cases in phase 2 had no relationship to the phase 1 cases in the alternate set.

In phase 2, all participants were asked to diagnose 8 new cases (the same for all participants), doing their best to provide an accurate diagnosis as quickly as possible. This procedure aimed at inducing nonanalytical reasoning based on pattern recognition, minimizing the chances that participants engage in elaborate analysis of case findings. The cases were presented in random order in a second booklet, and participants were reminded with each case to read the case description and then immedi-

ately write down the most likely diagnosis for the case. Four of the cases were similar to 2 cases seen in phase 1 by participants working with set 1, and the other 4 were similar to 2 cases seen in phase 1 by participants working with set 2 (Table 1). If the availability bias occurs, the diagnosis of the cases encountered in phase 1 should more promptly and frequently come to mind when participants encounter the cases with similar signs and symptoms in phase 2 than when they had not encountered these cases in phase 1. For example, participants working with set 1 in phase 1 would be expected to erroneously give a diagnosis of acute viral hepatitis to the cases of liver cirrhosis and primary sclerosing cholangitis more frequently than participants who worked with set 2 in phase 1.

In phase 3, participants were asked to again diagnose the 4 cases from phase 2 that could have been influenced by previous exposure to similar cases (Table 1). They followed instructions aimed at inducing reflective reasoning: (1) read the case; (2) write down the diagnosis previously given for the case; (3) list the findings in the case description that support this diagnosis; (4) list the findings that speak against this diagnosis; (5) list the findings that would be expected to be present if the diagnosis were true but were not described in the case. Participants were subsequently asked to list alternative diagnoses assuming that the initial diagnosis generated for the case had proved to be incorrect, and to follow the same procedure (steps 3-5) for each alternative diagnosis. Finally, they were asked to draw a conclusion by ranking the diagnoses in order of likelihood and selecting their final diagnosis for the case.

Data Analysis

All cases had a confirmed diagnosis that was used as a standard to evaluate the accuracy of the diagnoses provided by the participants. Two experts in internal medicine (J.L.C.M.S and C.G.) independently assessed the diagnoses blinded to the experimental conditions under which they were pro-

vided. The diagnoses were evaluated as fully correct, partially correct, or incorrect, and scored as 1, 0.5, or 0 points, respectively. A diagnosis was considered fully correct whenever the core diagnosis was cited by the participant and partially correct when the core diagnosis was not mentioned but a constituent element of the diagnosis was cited. For example, in the case in the Box, "celiac disease" was scored as correct, and "malabsorption" as partially correct.

For each participant, we separately summed the scores obtained in phase 2 on the 4 cases that had similarities to the cases encountered in phase 1 and the 4 cases that did not. For phase 3, the diagnostic scores obtained on the 4 cases were summed for each participant.

For phase 2, an analysis of variance (ANOVA) with years of training as a between-subjects factor (first vs second year) and recent experiences with similar cases as a within-subjects factor (with vs without) was conducted on the mean diagnostic performance scores obtained through nonanalytical reasoning on both types of cases (similar to cases seen in phase 1 or not). This analysis tested the hypothesis that recent experiences with similar cases would generate an availability bias and that this bias would be larger for more experienced, second-year residents. Post hoc paired *t* tests were performed to compare the diagnostic performance of first- and second-year residents under the 2 experimental conditions. To assess whether the diagnoses of the cases encountered in phase 1 were indeed provided as diagnosis of the similar cases in phase 2, we computed the number of times the diagnoses of cases in phase 1 were mentioned by participants in phase 2 who had seen similar cases in phase 1 vs those who had not and conducted paired *t* tests on these data for the first- and second-year residents.

A second ANOVA with years of training as a between-subjects factor (first year vs second year) and type of reasoning as a within-subjects factor (nonanalytical vs

Table 2. Mean Diagnostic Accuracy Scores in Phase 2 (Perfect Score, 4.0) as a Function of Previous Exposure to Similar Cases in Phase 1

	Training Year	
	First (n = 18)	Second (n = 18)
Diagnostic accuracy on the cases similar to those previously encountered	Mean Score (95% Confidence Interval) 2.03 (1.55-2.51)	1.55 (1.15-1.96)
Diagnostic accuracy on the other cases	1.42 (0.92-1.92)	2.19 (1.73-2.66)
<i>P</i> value	.046 ^a	.03 ^a

^aComparison of accuracy on similar vs other cases. Two-sided *t* test.

Table 3. Persistence of Phase 1 Diagnoses Among 8 Potentially Similar Cases in Phase 2 and 4 Similar Cases in Phase 3

	Training Year	
	First (n = 18)	Second (n = 18)
Frequency of phase 1 diagnosis given for phase 2 cases, mean (95% CI) [% of all wrong diagnoses]	Mean (95% CI) [% of All Wrong Diagnoses]	
Having encountered similar cases in phase 1	0.78 (0.34-1.26) [42]	1.44 (0.93-1.96) [63]
Not having encountered similar cases in phase 1	0.89 (0.47-1.30) [36]	0.72 (0.28-1.17) [42]
<i>P</i> value	.67 ^a	.04 ^a
	No. (%)	
Correction of phase 1 diagnoses in phase 3		
No. of phase 1 diagnoses given for phase 2 cases	14	26
Corrected after reflection in phase 3	5 (36)	8 (31)
Adhered to after reflection in phase 3	7 (50)	16 (63)
Changed to a different incorrect diagnosis	2 (14)	2 (6)

Abbreviation: CI, confidence interval.

^aComparison of phase 1 diagnoses with vs without having encountered similar cases in phase 1, 2-sided *t* test.

reflective) was conducted on the mean diagnostic performance scores in phase 2 and phase 3. This analysis tested the hypothesis that reflection (phase 3) could counteract the availability bias by improving the diagnostic performance scores compared with those obtained on the same cases through nonanalytical reasoning (phase 2).

Significance was set at $P < .05$ for all comparisons (2-tailed). SPSS version 15.0 (SPSS Inc, Chicago, Illinois) for Windows was used for the statistical analyses.

RESULTS

Nonanalytic Reasoning (Phase 2)

TABLE 2 presents the mean diagnostic accuracy scores obtained by first-year and second-year residents when cases were solved through nonanalytical reasoning (phase 2). The ANOVA showed no significant main effects, but there was a significant interaction effect between years of training and recent

experiences with similar cases ($F[1,34]=10.35$, mean square error (MSE)=0.68, $P=.003$, $\eta_p^2=0.23$). Mean scores for the second-year residents were consistent with an availability bias. They obtained significantly lower diagnostic scores on the cases similar to those encountered in phase 1 than the other cases (0-4 scale, 1.55; 95% confidence interval [CI], 1.15-1.96 vs 2.19; 95% CI, 1.73-2.66; $P=.03$).

Among the 8 phase 2 cases potentially similar to phase 1, second-year residents more frequently gave the phase 1 diagnosis when they had encountered the cases in phase 1 than when they had not (mean frequency per resident, 1.44; 95% CI, 0.93-1.96 vs 0.72; 95% CI, 0.28-1.17; $P=.04$) (TABLE 3). Even when the participants had not encountered the similar cases in phase 1, they sometimes incorrectly provided the phase 1 diagnosis to the related cases but this occurred less frequently than when they had been previously exposed to the phase 1 cases.

Table 4. Mean Diagnostic Accuracy Scores in Phase 2 (Perfect Score, 4.0) and Phase 3 for the Cases Similar to Those Encountered in Phase 1

	Training Year	
	First (n = 18)	Second (n = 18)
Nonanalytical diagnostic reasoning (Phase 2)	Mean Score (95% Confidence Interval) 2.03 (1.55-2.51)	1.55 (1.14-1.96)
Reflective diagnostic reasoning (Phase 3)	2.31 (1.89-2.73) ^a	2.03 (1.49-2.57) ^a

^aThere was a significant main effect of reflection, which improved diagnoses compared to nonanalytical reasoning for both 1st and 2nd year residents ($P = .006$ by ANOVA).

In contrast, this pattern was not seen for the first-year residents, who had a higher score on the cases similar to those encountered in phase 1 than on the other cases (Table 2). Having encountered a similar case in phase 1 did not lead to more frequently giving this diagnosis in phase 2 than when they had not seen a similar case (mean frequency per resident, 0.78; 95% CI, 0.34-1.26 vs 0.89; 95% CI, 0.47-1.30; $P = .67$) (Table 3).

Reflective Reasoning (Phase 3 vs Phase 2)

The diagnostic scores obtained through reflective reasoning (phase 3) on the cases similar to the diseases that had been encountered in phase 1 (those cases subject to an availability bias in phase 2) are presented in TABLE 4. A significant main effect of "type of reasoning" was found in the ANOVA ($F[1,34] = 8.46$, $MSE = 0.30$, $P = .006$, $\eta_p^2 = 0.20$) indicating that reflection improved all participants' diagnoses compared with nonanalytical reasoning. The percentage of phase 1 diagnoses that were corrected or adhered to after reflection is shown in Table 3.

COMMENT

This study demonstrated that an availability bias may indeed occur in response to recent experiences with similar clinical cases when a nonanalytical mode of reasoning is used, yielding diagnostic errors, and that reflective reasoning may help counteract this bias. The results suggest that the occurrence and negative effects of availability bias are a function of the reasoning approach and the expertise level.

Encountering only one case of a disease was sufficient to make the second-year residents more prone to incorrectly giving that diagnosis to subsequent cases of different, though similar, diseases. In emergency departments and outpatient clinics, physicians are likely to see (often close in time) several patients with similar symptoms caused by different diseases. In many clinical settings, therefore, conditions propitious for the occurrence of the availability bias prevail.

Moreover, because reliance on nonanalytical reasoning tends to increase with experience, it is possible that physicians with many years of clinical practice may be even more susceptible to availability bias than second-year residents, and this should be investigated. In real-life situations, an initial incorrect hypothesis might be spontaneously revised before expensive or time-consuming tests are ordered. However, the effects of anchoring by an early incorrect diagnosis may still lead to inaccurate judgment and inappropriate decisions. More experienced clinicians appear to be more subject to an anchoring effect,¹⁸ which makes it less likely that they will spontaneously overrule an incorrect initial diagnosis.

These findings contribute some insight into cognitive mechanisms underlying errors, which are the object of ongoing scientific debate.¹⁹ Evidence of the availability bias emerged in phase 2, when participants diagnosed the cases through a nonanalytical reasoning mode, and this was in part repaired in phase 3 by reflective reasoning. This suggests that mistakes made in phase 2 did not derive from lack of knowledge. Residents who failed to correctly diagnose the cases through

nonanalytical reasoning may have arrived at the correct diagnoses after reflecting on the same cases by activating existing knowledge. Therefore, errors in phase 2 were more likely to have been provoked by bias in the reasoning processes.

We had expected the availability bias to be larger for the more experienced residents because the tendency to diagnose cases through pattern recognition increases with clinical experience.^{12,13} However, we had not expected to find an opposite pattern for the first-year residents, who had better performance on similar cases. It is possible to speculate on reasons for this finding, such as that these novice residents might have already used a more reflective mode of reasoning during the exposure phase (phase 1), being less self-confident than their more experienced colleagues^{20,21} and, therefore, perhaps less reliant on immediate decisions. They may not have had a sufficient amount of clinical experience to make extensive use of pattern recognition and had to rely on a more analytic approach that could have been activated by phase 1 cases. However, as a post hoc analysis yielding an unexpected finding, these are speculations that should only be interpreted as hypothesis generating.

Although reliance on nonanalytical reasoning and heuristics such as availability work well in many situations, reducing the time and effort involved in decision making and allowing physicians to make accurate diagnoses in routine situations,^{19,22} it may open the door to cognitive bias. Reflection has been shown to improve diagnosis when problems are complex or nonroutine,^{17,23} and this study indicates that reflection may also be a mechanism to counteract cognitive biases.

With respect to medical education, this study suggests that a relatively simple instructional procedure can be used to induce reflective reasoning and improve diagnostic accuracy. This procedure for reflective reasoning can be implemented relatively easily in educational situations. Further research should investigate the effects

of this process on diagnostic reasoning in practice settings.

This study has several limitations. First, we investigated residents from 2 different years in the internal medicine residency program, and it is not clear whether the differences in the susceptibility to bias encountered in the study would persist in later years or occur in other specialties. Second, the test cases were presented immediately after the initial cases and similar problems do not always come consecutively in real clinical practice. Third, there may be restrictions in generalizing these findings obtained under laboratory conditions to real-life situations, which are always richer in

cues that may facilitate intuitive judgments. However, we worked with cases based on real patients and with tasks that simulate medical decision making.

In summary, this study showed that the availability bias may occur in medical diagnosis as a consequence of recent experiences with similar cases under nonanalytical reasoning conditions and that susceptibility to this effect may be related to having more clinical experience. It provided further evidence that flaws in reasoning processes rather than knowledge gaps may underlie diagnostic errors and showed the potential for repair by reflective reasoning.

Author Contributions: Dr Mamede had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Mamede, Berge, Rikers, Schmidt.

Acquisition of data: Berge, Saase, Guldener.

Analysis and interpretation of data: Mamede, Gog, Berge, Rikers, Saase, Guldener, Schmidt.

Drafting of the manuscript: Mamede, Gog.

Critical revision of the manuscript for important intellectual content: Mamede, Gog, Berge, Rikers, Saase, Guldener, Schmidt.

Statistical analysis: Mamede, Gog.

Administrative, technical, or material support: Rikers, Saase, Guldener.

Study supervision: Schmidt.

Financial Disclosures: None reported.

Funding/Support: None.

Additional Contributions: We thank Júlio César Penaforte, MD, MSc (Hospital Geral de Fortaleza, Brazil) and João Macedo Coelho Filho, MD, PhD, (Faculty of Medicine, Federal University of Ceará, Brazil) for their permission to use the clinical cases that they prepared for previous studies, without compensation.

REFERENCES

- Graber ML, Franklin N, Gordon R. Diagnostic error in internal medicine. *Arch Intern Med*. 2005;165(13):1493-1499.
- Kohn KT, Corrigan JM, Donaldson MS. *To Err Is Human: Building a Safer Health System*. Washington, DC: National Academy Press; 1999.
- Kahnemann D, Slovic P, Tversky A. *Judgment Under Uncertainty: Heuristics and Biases*. New York, NY: Cambridge University Press; 1982.
- Croskerry P. The importance of cognitive errors in diagnosis and strategies to minimize them. *Acad Med*. 2003;78(8):775-780.
- Kahneman D. A perspective on judgment and choice: mapping bounded rationality. *Am Psychol*. 2003;58(9):697-720.
- Klein JG. Five pitfalls in decisions about diagnosis and prescribing. *BMJ*. 2005;330(7494):781-783.
- Redelmeier DA. The cognitive psychology of missed diagnoses. *Ann Intern Med*. 2005;142(2):115-120.
- Poses RM, Anthony M. Availability, wishful thinking, and physicians' diagnostic judgments for patients with suspected bacteremia. *Med Decis Making*. 1991;11(3):159-168.
- Brezis M, Halpern-Reichert D, Schwaber MJ. Mass media-induced availability bias in the clinical suspicion of West Nile fever. *Ann Intern Med*. 2004;140(3):234-235.
- Heath L, Acklin M, Wiley K. Cognitive heuristics and AIDS risk assessment among physicians. *J Appl Soc Psychol*. 1991;21:1859-1867.
- Peay MY, Peay ER. The evaluation of medical symptoms by patients and doctors. *J Behav Med*. 1998;21(1):57-81.
- Schmidt HG, Boshuizen HP. On acquiring expertise in medicine. *Educ Psychol Rev*. 1993;5:1-17.
- Schmidt HG, Norman GR, Boshuizen HP. A cognitive perspective on medical expertise: theory and implication. *Acad Med*. 1990;65(10):611-621.
- Evans JS. Dual-processing accounts of reasoning, judgment, and social cognition. *Annu Rev Psychol*. 2008;59:255-278.
- Eva KW. What every teacher needs to know about clinical reasoning. *Med Educ*. 2005;39(1):98-106.
- Mamede S, Schmidt HG, Rikers RM, Penaforte JC, Coelho-Filho JM. Breaking down automaticity: case ambiguity and shift to reflective approaches in clinical reasoning. *Med Educ*. 2007;41(12):1185-1192.
- Mamede S, Schmidt HG, Penaforte JC. Effects of reflective practice on the accuracy of medical diagnoses. *Med Educ*. 2008;42(5):468-475.
- Eva KW. The aging physician: changes in cognitive processing and their impact on medical practice. *Acad Med*. 2002;77(10)(suppl):S1-S6.
- Norman G, Eva KW. Diagnostic error and clinical reasoning. *Med Educ*. 2010;44(1):94-100.
- Friedman CP, Gatti GG, Franz TM, et al. Do physicians know when their diagnoses are correct? implications for decision support and error reduction. *J Gen Intern Med*. 2005;20(4):334-339.
- Berner ES, Maisiak RS, Heudebert GR, Young KR Jr. Clinician performance and prominence of diagnoses displayed by a clinical diagnostic decision support system. *AMIA Annu Symp Proc*. 2003;76-80.
- Norman G. Dual processing and diagnostic errors. *Adv Health Sci Educ Theory Pract*. 2009;14(suppl 1):37-49.
- Mamede S, Schmidt HG, Rikers RM, Penaforte JC, Coelho-Filho JM. Influence of perceived difficulty of cases on physicians' diagnostic reasoning. *Acad Med*. 2008;83(12):1210-1216.