
# **Cochlear Implants**

ochlear implants are small, electronic devices that can be surgically implanted to enable profoundly deaf children and adults who are not helped by traditional hearing aids to hear better. The May 19, 2004, issue of *JAMA* includes an article about cochlear implantation in children, and the May 2004 issue of the *Archives of Otolaryngology–Head* & Neck Surgery is a theme issue devoted to this topic.

## HEARING

- Sound waves funneled into the ear cause the tympanic membrane (eardrum) to vibrate.
- These vibrations are transmitted across a bridge formed by 3 small bones to the cochlea of the inner ear.
- The fluid-filled cochlea contains hair cells that sense vibrations transmitted through the fluid.
- The hair cells trigger impulses in the auditory (hearing) nerve that transmit to the brain where they are interpreted as sound.

Most profound deafness is **sensorineural**—there is damage to the sensitive and vulnerable hair cells in the cochlea so that vibrations in the cochlea cannot be transmitted. Cochlear implants can restore hearing by bypassing the hair cells and stimulating the auditory nerve



## FOR MORE INFORMATION

- American Speech-Language-Hearing Association
   1-800/638-8255
   TTY: 1-800/498-2071
   www.asha.org
- Cochlear Implant Association, Inc www.cici.org
- Self-Help for Hard of Hearing People Voice: 301/657-2248 TTY: 301/657-2249 www.shhh.org

#### **INFORM YOURSELF**

To find this and other JAMA Patient Pages, go to the Patient Page link on JAMA's Web site at www.jama.com. A Patient Page on adult hearing loss was published in the April 16, 2003, issue.

Sources: Cochlear Implant Association, Inc, National Institute on Deafness and Other Communication Disorders, American Speech-Language-Hearing Association

#### **AFTER SURGERY**

- After the patient recovers from surgery for about a month, the speech processor is placed and then adjusted as hearing improves in a process called **mapping**. As hearing gets better, further adjustments are made over time.
- Training with speech therapists or a teacher of the hearing-impaired will help patients learn how to use the cochlear implant. Children in their formative years need to be taught to understand new sounds and their meanings and to translate them into speech and language.

Sharon Parmet, MS, Writer
Cassio Lynm, MA, Illustrator
Richard M. Glass, MD, Editor

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. Any other print or online reproduction is subject to AMA approval. To purchase bulk reprints, call 718/946-7424.

