

Review

Sigmoid Diverticulitis

A Systematic Review

Arden M. Morris, MD, MPH; Scott E. Regenbogen, MD, MPH; Karin M. Hardiman, MD, PhD;
Samantha Hendren, MD, MPH

IMPORTANCE Diverticulitis is a common disease. Recent changes in understanding its natural history have substantially modified treatment paradigms.

OBJECTIVE To review the etiology and natural history of diverticulitis and recent changes in treatment guidelines.

EVIDENCE REVIEW We searched the MEDLINE and Cochrane databases for English-language articles pertaining to diagnosis and management of diverticulitis published between January 1, 2000, and March 31, 2013. Search terms applied to 4 thematic topics: *pathophysiology*, *natural history*, *medical management*, and *indications for surgery*. We excluded small case series and articles based on data accrued prior to 2000. We hand searched the bibliographies of included studies, yielding a total of 186 articles for full review. We graded the level of evidence and classified recommendations by size of treatment effect, according to the guidelines from the American Heart Association Task Force on Practice Guidelines.

FINDINGS Eighty articles met criteria for analysis. The pathophysiology of diverticulitis is associated with altered gut motility, increased luminal pressure, and a disordered colonic microenvironment. Several studies examined histologic commonalities with inflammatory bowel disease and irritable bowel syndrome but were focused on associative rather than causal pathways. The natural history of uncomplicated diverticulitis is often benign. For example, in a cohort study of 2366 of 3165 patients hospitalized for acute diverticulitis and followed up for 8.9 years, only 13.3% of patients had a recurrence and 3.9%, a second recurrence. In contrast to what was previously thought, the risk of septic peritonitis is reduced and not increased with each recurrence. Patient-reported outcomes studies show 20% to 35% of patients managed nonoperatively progress to chronic abdominal pain compared with 5% to 25% of patients treated operatively. Randomized trials and cohort studies have shown that antibiotics and fiber were not as beneficial as previously thought and that mesalamine might be useful. Surgical therapy for chronic disease is not always warranted.

CONCLUSIONS AND RELEVANCE Recent studies demonstrate a lesser role for aggressive antibiotic or surgical intervention for chronic or recurrent diverticulitis than was previously thought necessary.

+ Supplemental content at
jama.com

+ Related article at
jamasurgery.com

+ CME Quiz at
jamanetworkcme.com and
CME Questions 300

Author Affiliations: Division of Colorectal Surgery, Department of Surgery, University of Michigan, Ann Arbor.

Corresponding Author: Arden M. Morris, MD, MPH, Division of Colorectal Surgery, Department of Surgery, University of Michigan, 1500 E Medical Center Dr, TC-2124, Ann Arbor, MI 48109-5343 (ammsurg@med.umich.edu).

Section Editor: Mary McGrae McDermott, MD, Senior Editor.

JAMA. 2014;311(3):287-297. doi:10.1001/jama.2013.282025

Before effective broad-spectrum antibiotics were available, diverticulitis was a devastating disease associated with substantial morbidity and mortality. Now, most cases resolve with antibiotic therapy. Because of the fear of complications associated with perforation, if multiply recurrent disease occurs, surgical resection of the involved colon is performed.¹ However, recent advances in the understanding of the disease's pathophysiology and natural history have led to substantial changes in diverticulitis treatment guidelines.

Over the past decade, 4 key innovations have changed the thinking about and management of diverticulitis: complicated diverticulitis (ie, with perforation, abscess, or phlegmon) is now reliably distinguished from uncomplicated disease by computed tomography (CT)²; large clinical and administrative databases have facilitated more complete follow-up of large populations, resulting in changes in the understanding of the natural history of diverticulitis, clinical and behavioral risk factors for the disease, and what the indications and outcomes of its treatments are; similarities exist between the physiology and inflammatory processes for diverticulitis, irritable bowel syndrome, and inflammatory bowel disease. These insights led to new approaches to managing chronic disease and preventing recurrent diverticulitis.³ Surgeons now pursue less invasive intervention, increasing the use of percutaneous drainage, intraperitoneal lavage, and minimally invasive surgical techniques.⁴

Diverticulitis treatments are rapidly evolving. This review summarizes recent medical literature describing the pathophysiology and natural history of chronic and recurrent sigmoid diverticulitis and reviews new recommendations for the medical management and indications for surgery. Four key questions are addressed: (1) What is known about the pathophysiology of diverticulitis? How do diverticula become inflamed, and what are associated risk factors? (2) How have large observational trials clarified the natural history of diverticulitis that is managed nonoperatively? (3) What are the proposed mechanisms, options, and outcomes of medical therapy for diverticulitis? and (4) What are the indications for surgical treatment of diverticulitis?

Methods

Data Sources

We performed a systematic review of the MEDLINE and Cochrane databases, using separate search terms for each of the 4 key questions (eAppendix in the Supplement), for articles published between January 1, 2000, and March 31, 2013. Broad search terms for question 1 included (*pathophysiology or etiology or pathogenesis*) and *diverticulitis*; for question 2, broad search terms were (*natural history or outcome*) and *diverticulitis*; for questions 3 and 4, broad search terms were (*chronic or acute or smoldering or recurrent*) and *diverticulitis* and (*management or treatment*). Searches were limited to English language articles published since 2000 that addressed diverticulitis in adult humans. All articles were then combined into a single list, and duplicates were excluded, resulting in 1383 abstracts and articles for review.

Study Selection

We reviewed abstracts and excluded commentary or opinion pieces, review articles that reported data present in other included refer-

ences, articles based on data accrued before 2000, and articles containing primary data duplicated in another included article. In the case of duplicate presentation of data, we selected articles with the most recent analyses. We excluded small case series with fewer than 30 patients, except for the search of articles for question 1 regarding pathophysiology. We supplemented our automated search by manually searching additional references from the bibliographies of included studies, yielding a total of 186 articles for full review.

Data Extraction

Studies selected for inclusion were reviewed according to guidelines from the Strengthening the Reporting of Observational Studies in Epidemiology.⁵ We graded the level of evidence and classified recommendations by size of treatment effect, according to the guidelines from the American Heart Association Task Force on Practice Guidelines.⁶ We compared the resulting recommendations with those of the most comprehensive recent clinical practice guidelines on management of diverticulitis.⁴

Results

Eighty articles were selected from the 186 reviewed. Articles were excluded if there was an absence of data regarding the topic of interest, if the majority of data accrued prior to 2000, and if it was a small case series. Articles were organized into 4 categories consistent with the study questions and summarized below.

Pathophysiology and Risk Factors

Twenty-five articles fulfilled criteria and were included in this review of the pathophysiology of diverticulitis. Although *colonic diverticulum* refers specifically to a thin-walled outpouching of the mucosa and serosa, absent the muscularis, and *diverticulosis* refers to the presence of many diverticula, *diverticulitis* is distinguished by the presence of inflammation. If untreated, diverticular inflammation may resolve, become chronic, or progress, leading to bacterial translocation or even perforation of the colon wall at the inflamed site.

The prevailing explanation for colonic diverticula formation posits that altered bowel motility leading to increased intraluminal pressure causes mucosal outpouching adjacent to the vasa recta. Whether diverticula, once formed, can spontaneously resolve is unknown. The mechanism by which asymptomatic diverticula become inflamed and perforate (diverticulitis) is still under investigation but is plausibly associated with altered gut motility and increased pressure combined with a deranged colonic microenvironment. In recent series, when ex vivo colonic tissue from patients with diverticulitis was exposed to chemicals that contract or relax smooth muscle, the response was significantly abnormal with increased hypercontractility and lower maximum relaxation responses.⁷⁻⁹ These findings are consistent with neuropeptide abnormalities and the altered histologic appearance of muscle and nerves in the bowel wall of patients with diverticulitis compared with healthy patients. For example, reduced serotonin transporter expression and fewer interstitial cells of Cajal were found among patients with diverticulitis but not among those with normal colons or with noninflamed diverticula.¹⁰⁻¹² Moreover, patients who ingest calcium channel blockers, which reduce smooth muscle contractility, appear to have a re-

duced risk of perforated diverticulitis compared with patients who do not take calcium blockers.¹³

Recurrent or chronic diverticulitis displays chemical and histological similarities to inflammatory bowel disease and irritable bowel syndrome.^{14,15} These diagnoses may be concurrent, sequential, or entirely separate. Although the mechanisms of inflammation are unknown, higher levels of histamine, tumor necrosis factor α (TNF- α), and matrix metalloproteinases have been identified in colonic biopsies from patients with irritable bowel syndrome, inflammatory bowel disease, and diverticulitis.¹⁶⁻¹⁸ Other common evidence of chronic inflammation includes the presence of granulomas and infiltrating lymphocytes. Ultimately, however, there were minimal mechanistic data to support or refute a common or related pathway and key distinguishing features of each persist. For example, a hallmark of *irritable bowel syndrome* is relief of crampy pain upon defecation; *inflammatory bowel disease* is characterized by mucosal injury frequently resulting in bloody diarrhea; *diverticulitis* is by definition associated with diverticula.

Lifestyle risk factors such as diet, smoking, and medication use have long been considered important in the etiology of diverticular disease. A prospective UK population-based cohort study found a relative risk of 0.69 (95% CI, 0.55-0.86) of diverticular disease among vegetarians compared with meat eaters.¹⁹ The relationship between dietary fiber and diverticula is not clear, however. A large cohort study of patients undergoing colonoscopy found that those who reported the highest fiber intake were at highest risk of diverticulosis.²⁰ A longitudinal survey of 47 228 health professionals recently reported that incident diverticulitis was not associated with nut, corn, or popcorn ingestion and that increased nut intake was associated with lower risk of diverticulitis.²¹ Smoking and obesity have been linked to diverticulitis and to complicated diverticulitis in several large cohort studies,²²⁻²⁶ whereas increased physical activity is associated with decreased risk.^{27,28} Nonsteroidal anti-inflammatory drugs, opioids, and corticosteroids have been convincingly associated with increased risk of perforated diverticulitis.²⁹⁻³² A unifying hypothesis to integrate these diverse lifestyle effects and their contribution to pathophysiology has not been advanced.

Natural History

We defined the *natural history of diverticulitis* as the longitudinal outcomes for patients whose disease was managed nonoperatively. Six articles fulfilled criteria and were reviewed. Most of the data regarding natural history were focused on nonoperative outcomes and specifically on risk of recurrence, that is, subsequent acute diverticulitis after an asymptomatic interval, rather than on chronic disease in which there is no asymptomatic interval.

Risk of Recurrence

Two large multicenter studies^{33,34} confirmed that recurrence is rare and is a relatively benign process for the substantial majority of patients. Broderick-Villa et al³⁴ reported on 2366 of 3165 patients (75%) hospitalized with acute diverticulitis and treated nonoperatively in the Kaiser Permanente system. Eighty-six percent of those patients required no further inpatient care for diverticulitis over the 8.9 years of follow-up. Recurrence occurred in only 13.3% of patients and only 3.9% had a second recurrence. No patient with a second recurrence required an operation, and repeat recurrences plateaued

after 4 episodes. Although the risk for a second recurrence increased to 29% among those with a first recurrence, the authors concluded that recurrence overall is rare and therefore does not warrant elective colectomy.

Binda et al³³ obtained complete follow-up for 320 patients treated with antibiotics in 17 Italian hospitals after admission for acute diverticulitis. Over a mean period of 10.7 years, 61% of patients required no further inpatient care. Twenty-two percent of patients had persistent or recurrent symptoms requiring hospitalization, and 17% had a recurrent episode resulting in an emergency operation. In adjusted analyses, the risk of recurrence was greatest among patients younger than 50 years and among those with at least 3 previous episodes. The authors noted that unlike other studies of recurrence, episodes were not clustered into the first 2 years or even the first 5 years following the index diagnosis.

Complications of Diverticulitis

A large population-based study linked primary care and hospital-based data to examine patterns of morbidity associated with complicated diverticulitis among 2950 patients hospitalized in the United Kingdom.³⁵ Seventy-two percent of patients had no antecedent episodes of diverticulitis, and 2 or more prior episodes were not associated with abscess or stricture formation, although they were associated with increased risk of fistula. Although most serious complications of diverticulitis were linked to the first episode, not to recurrence, they were consequential. Compared with age- and sex-matched cases in the general population, patients with perforation or abscess had 4.5-fold increased risk of death in the ensuing year.

Shifting Morphology

Morphologic characteristics of recurrent diverticulitis may differ considerably from the first episode. A single institution study³⁶ of 60 patients with recurrent disease after initial medical management of uncomplicated diverticulitis compared CT scans from the index and second admission. At the time of recurrence, 6 patients (10%) had complicated diverticulitis and 3 underwent an urgent Hartmann procedure. Among the 54 patients with a CT-defined recurrence that was uncomplicated, 19 (35%) had a morphologically distinct recurrence of diverticulitis at an average of 8 cm from the previous site.

Chronic Pain

Nelson et al³⁷ analyzed a cohort of 99 patients with complicated diverticulitis whose care was managed nonoperatively with follow-up over 76 months. Forty-six patients had recurrent disease and 20 underwent an elective resection more than 6 months after the initial episode. Unlike Binda et al, the authors found no difference between recurrence or need for emergency operation based on age. Although about half of the patients had recurrent or chronic disease, none died of complications of diverticulitis. Similarly, a longitudinal survey of 124 patients with diverticular disease who were managed medically over 7 years reported that 34% continued to experience abdominal pain for 3.5 days per month. Odds of chronic pain were increased 4-fold among those who had been previously diagnosed with diverticulitis.³⁸

Until recently, recommendations for management and prevention of diverticulitis were well-established. Patients with perforation, abscess, fistula, or stricture were managed definitively with an

Table 1. Medical Management of Chronic and Recurrent Diverticulitis

| Source | Evidence | | Sample Size ^c | Study Design | Intervention | Primary Outcome(s) | Main Finding(s) | P Value |
|---------------------------------------|--------------------|--------------------|--------------------------|-----------------------------------|---|---------------------------------------|---|---------|
| | Level ^a | Class ^b | | | | | | |
| Alonso et al, ³⁹ 2010 | B | IIb | 70 | Cohort | 7-d Oral antibiotics | Symptom relief | 97% Outpatient symptom relief | |
| Bianchi et al, ⁴⁰ 2011 | C | IIb | 1660 | Meta-analysis | Rifaximin + fiber vs fiber | 1-y Symptom relief | 29% Rifaximin + fiber relief | .001 |
| Brandimarte et al, ⁴¹ 2004 | C | IIa | 86 | Cohort | Mesalamine and rifaximin, then 8-wk mesalamine | Symptom relief | 78% Asymptomatic | |
| Chabok et al, ⁴² 2012 | A | III | 623 | RCT | Antibiotics vs none | Recurrence prevention | 16 % Recurrence in both groups | .88 |
| Comparato et al, ⁴³ 2007 | B | IIa | 244 | RCT | High- vs low-dose rifaximin vs high- vs low-dose mesalamine | Symptom relief | Mesalamine > rifaximin high dose > low dose | |
| De Korte et al, ⁴⁴ 2011 | B | III | 3 Studies | Retrospective cohort study + RCTs | Antibiotics vs none | Symptom relief | No significant difference | |
| Dughera et al, ⁴⁵ 2004 | B | IIa | 76 | RCT | Oral probiotics vs placebo | Symptom relief, recurrence prevention | Reduced symptoms with treatment, but recurrence no difference | .28 |
| Hjern et al, ⁴⁶ 2007 | B | III | 311 | Retrospective cohort | Antibiotics vs none | Symptom relief | 29% Recurrence in treated vs 28% in not treated | |
| Mizuki et al, ⁴⁷ 2005 | B | IIa | 65 | Cohort | 10-d oral antibiotics and gradual diet resumption | Recurrence prevention | 25% Recurrence | |
| Moya et al, ⁴⁸ 2012 | C | III | 76 | Cohort | IV vs oral antibiotics | Symptom relief, cost | No significant difference | |
| Ribas et al, ⁴⁹ 2010 | B | IIb | 50 | RCT | Short vs long course | Symptom relief | No significant difference | |
| Ridgeway et al, ⁵⁰ 2009 | B | IIa | 79 | RCT | Oral vs IV antibiotics | Symptom relief | No significant difference | |
| Shabanzadeh et al, ⁵¹ 2012 | B | III | 3 Studies | Review | Antibiotics vs none | Recurrence prevention | No significant difference | |
| Tursi et al, ⁵² 2002 | B | IIb | 193 | RCT | Cyclic combined rifaximin and mesalamine vs cyclic rifaximin only | Recurrence prevention | 3% Recurrence combined therapy vs 12% rifaximin only | .005 |

Abbreviations: IV, intravenous; RCT, randomized clinical trial.

^a Level of evidence indicates the precision of the estimate of treatment effect.

^b Class of recommendation indicates the size of the treatment effect. Level A indicates the strongest weight of evidence; Level B is intermediate; Level C

indicates the lowest rank of evidence, for example, when evidence is based on expert consensus.

^c Sample size indicates the number of study patients for whom complete data were available.

urgent or elective sigmoid colon resection. Those with uncomplicated disease were managed with antibiotics and bowel rest. In the event of a recurrence or failure to resolve, patients were scheduled for an elective sigmoid colectomy. The rationale for elective surgery was largely preventive, based on concerns that recurrence would result in progressively increased risk of sepsis or the need for a colostomy. More recently, emerging medical therapies are under investigation as a potentially lower-risk means of prevention.

Emerging Medical Therapies

The goals of medical therapy for diverticulitis are to decrease inflammation acutely, to prevent recurrence, and to manage chronic symptoms. With the advantage of more current pathophysiologic data and large database-derived studies of natural history, new medical approaches toward these goals have been proposed. A total of 14 articles on existent or emerging medical therapies met criteria for review (Table 1); however, the quality of most available evidence was poor (level B or C) and only 1 article met level A criteria.

Fiber


The traditional approach to prevention of recurrence of diverticulitis has been to increase dietary or supplemental fiber. Ünlü et al⁵³

recently published a systematic review of the evidence supporting this longstanding recommendation. The authors found 4 studies on treatment and no studies on prevention of recurrence that met inclusion criteria. One randomized trial showed no treatment effect of fiber ingestion on resolution of symptoms, whereas the other 3 studies did show a significant treatment advantage with fiber ingestion. The authors concluded that recommendations for ingestion for dietary fiber are based on inconsistent level 2 and level 3 evidence. None of the studies examined met criteria for inclusion in our study because the data were collected prior to 2000. A systematic review⁴⁰ of 4 trials of fiber vs rifaximin plus fiber found that the combined therapy was slightly but significantly more effective in obtaining symptom relief and preventing complications at 1 year. Two of the 4 studies were based on data from prior to 2000, the third was not explicitly about diverticulitis, and the fourth did not state what years data were collected.

Antibiotic Therapy

A major change in the approach to the management of acute uncomplicated diverticulitis is the progressively reduced use of antibiotics. Specifically, prospective randomized and open trials have shown no advantage of intravenous over oral antibiotics, and there-

Figure. Clinical Outcomes Based on Current Treatment Standards for a Hypothetical Cohort of 1000 Patients Presenting With Acute Diverticulitis



Data derived from the following studies of diverticulitis outcomes: Ambrosetti et al,⁵⁹ Broderick-Villa et al,³⁴ Nelson et al,³⁷ Kaiser et al,⁶⁰ Dharmarajan et al,⁵⁷ Anaya et al,⁷¹ and Hall et al.⁷⁰

^aComplicated diverticulitis refers to the presence of perforation, abscess, or phlegmon.

fore have recommended outpatient management.^{39,47-50} In addition, more recent data indicate equivalent efficacy of a 4-day vs 7-day course of an appropriately broad-spectrum antibiotic.⁵⁴ This study recommended short-course therapy with ertapenem but rifaximin has also been favored for chronic or recurrent diverticulitis due to limited gut absorption and low cost.⁵² A Cochrane review⁵¹ of antibiotic use in acute uncomplicated diverticulitis found that, in spite of published guidelines, the best available data⁴² do not support use of antibiotics. A more inclusive systematic review⁴⁴ and a retrospective cohort study⁴⁶ have also found that antibiotic use has no effect on complications, need for surgery, or recurrence rate. Thus, newer data support a noninterventional policy for treatment of uncomplicated diverticulitis.

Probiotics

The rationale for use of probiotics in diverticular disease is based on the theory that a deranged colonic microenvironment, including abnormal gut flora, precipitates chronic inflammation and recurrent disease. One study of probiotics met inclusion criteria.⁴⁵ The authors randomized 83 consecutive patients, whose index episode was resolved after treatment with rifaximin or ciprofloxacin, to receive an oral polybacterial lysate suspension or placebo twice daily for 2 weeks every month within 3 months after an acute attack. Complete follow-up data were available for 76 patients. The probiotic group reported significantly less abdominal pain, bloating, and fever, but there was no significant difference in recurrence rates (2 of 41 vs 5 of 35).

Anti-inflammatory Medication

A recent appreciation of chronic mucosal inflammation has sparked interest in alterations of the colonic microenvironment and the potential for use of anti-inflammatory medication. A head-to-head comparison of anti-inflammatory treatment using mesalamine vs anti-

biotic therapy with rifaximin demonstrated significantly reduced symptoms after 6 to 12 months of high-dose cyclic mesalamine.⁴³ A nonrandomized comparison⁵² of rifaximin and mesalazine vs rifaximin alone showed 3% recurrence in the combined medication group and 13% recurrence in the rifaximin alone group over 1 year. A cohort study from the same authors⁴¹ showed that combined mesalamine and rifaximin followed by mesalamine alone led to resolution of symptoms for nearly all patients but recurrent disease among 2% at 8 weeks. The duration of follow-up for this study was notably short; most studies of recurrence require a minimum asymptomatic period of 3 to 6 months from the time of the index diagnosis before the next symptom onset.

Current Indications for Surgical Treatment

Surgery for acute diverticulitis is indicated for patients who present with sepsis and diffuse peritonitis and for patients whose condition did not improve with medical therapy, percutaneous drainage, or both.^{4,55} Surgical options include simple colostomy formation in the setting of profound inflammation, traditional sigmoid resection with colostomy (Hartmann procedure), and sigmoid resection with a primary colocolonic or colorectal anastomosis with or without a diverting loop ileostomy. Based on selection criteria, we identified 35 relevant articles describing indications for surgical therapy (in the Supplement). Most articles met only level B or C criteria and none met level A criteria, thereby limiting the conclusions that can be drawn.

Urgent Setting

In 2 cohort studies, the presence of complicated diverticulitis, abscess, free intraperitoneal air on computed tomography alone, or all 3 did not mandate an urgent operation among hemodynamically stable patients.^{56,57} Still, up to 25% of patients with evidence of abscess, perforation, or both underwent surgery during the acute in-

Table 2. Level of Recommendation for Systematic Review of Recent Literature Compared to Current Practice Guidelines for Prevention of Recurrent Sigmoid Diverticulitis

| Intervention | Current Evidence Review and Guidelines ^a | Recommendation ^b | |
|---|---|-----------------------------|-------|
| | | Level | Class |
| Recovered From 1 or More Uncomplicated Episode | | | |
| Fiber supplementation | | C | IIa |
| Evidence review | Not addressed. | | |
| Practice guidelines ^c | Long-term fiber supplementation may prevent recurrence (ASCRS) | | |
| Antibiotic use | | A | III |
| Evidence review | For acute uncomplicated diverticulitis, a Cochrane review, ^{42,51} a systematic review, ⁴⁴ and a retrospective cohort study ⁴⁶ do not support use of antibiotics for prevention of recurrence | | |
| Practice guidelines ^c | Not addressed | | |
| Probiotics | | C | IIb |
| Evidence review | A trial of 83 patients randomized to receive oral polybacterial lysate vs placebo reported no significant difference in recurrence rates ($P = .2$ using χ^2 comparison of proportions) ⁴⁵ | | |
| Practice guidelines ^c | Not addressed | | |
| Mesalamine | | B | IIa |
| Evidence review | Small uncontrolled trials indicate approximately 3% recurrence rate over 1 y with use of combined mesalamine and rifaximin ⁵² | | |
| Practice guidelines ^c | Not addressed | | |
| Avoiding nuts and seeds | | A | III |
| Evidence review | A survey of 47 228 health professionals reported that incident diverticulitis was not associated with nut, corn, or popcorn ingestion and that increased nut intake was associated with lower risk of diverticulitis ²¹ | | |
| Practice guidelines ^c | Not addressed | | |
| Surgical resection | | B | IIb |
| Evidence review | Several cohort studies compared recurrence among patients who treated operatively vs nonoperatively. They noted that a complicated recurrence occurred in fewer than 5% of patients treated nonoperatively ^{62,69-73} ; the occurrence of multiple subsequent episodes did not increase the risk of major complications of diverticulitis ⁷⁴ ; and complicated diverticulitis most commonly occurred during the first episode rather than during recurrent episodes. ⁷⁵⁻⁷⁸ Taken together, these data support a real but limited role for surgery in preventing recurrence of diverticulitis | | |
| Practice guidelines ^c | The decision to recommend elective sigmoid colectomy after recovery from acute diverticulitis should be made on a case-by-case basis (ASCRS) | B | I |
| | Indications for surgery most frequently reported include: ≥ 2 episodes of diverticulitis severe enough to cause hospitalization and any episode of diverticulitis associated with contrast leakage, obstructive symptoms, or an inability to differentiate between diverticulitis and cancer (WGO) | C | IIb |
| | Elective sigmoid resection may not be necessary after any specific number of episodes of uncomplicated diverticulitis or with any definite age thresholds (SSAT) | C | IIb |
| | Recurrent diverticulitis is rare after surgery (1%-10%) (WGO) | C | IIb |
| | Extraluminal air, contrast or abscess is a predictor of an adverse natural history and may be helpful in determining the need for surgery (ASCRS) | | |

(continued)

dex hospitalization.⁵⁸⁻⁶⁰ Elective surgery after successful nonoperative management of an episode of complicated diverticulitis was often recommended due to rates of recurrence, hospital readmission, and need for elective resection as high as 50 to 70,^{34,58-61} especially among those with a pelvic or paracolic abscess that required percutaneous drainage.⁵⁹ However, few of these patients needed an emergency operation.^{37,62}

Preoperative Colonoscopy

Elective evaluation of patients who recovered from an episode of acute diverticulitis was controversial. Some authors advocated colo-

noscopy to confirm the diagnosis and exclude malignancy.^{63,64} Others reported no increased detection of advanced neoplasia in patients with a typical presentation of acute diverticulitis.⁶⁵⁻⁶⁸

Recurrent Diverticulitis

Among studies comparing surgical with nonsurgical management, several important considerations challenged routine elective surgical therapy for recurrent or chronic diverticulitis. First, complicated recurrence after recovery from an uncomplicated episode of diverticulitis occurred in fewer than 5% of patients whose care was managed nonoperatively.^{62,69-73} Second, the occurrence of multiple sub-

Table 2. Level of Recommendation for Systematic Review of Recent Literature Compared to Current Practice Guidelines for Prevention of Recurrent Sigmoid Diverticulitis (continued)

| Intervention | Current Evidence Review and Guidelines ^a | Recommendation ^b | |
|-----------------------------------|--|-----------------------------|-------|
| | | Level | Class |
| Young Patients (≤50 y) | | | |
| Surgical resection | | C | IIb |
| Evidence review | Several cohort studies found modestly higher rates of recurrence among patients younger than 40 y than among those older than 40 y ^{71,72,85,86} ; however, these data were countered by other cohort studies that did not document a more aggressive disease course based on age ⁸⁷⁻⁹³ | | |
| Practice guidelines ^c | The decision to recommend elective sigmoid colectomy after recovery from acute diverticulitis should be made on a case-by-case basis (ASCRS) | B | I |
| | There is no clear consensus regarding whether younger patients (<50 y) are at increased risk of complications; however, they are probably at increased risk of recurrent diverticulitis (ASCRS) | C | IIa |
| | In young patients with no comorbid conditions, elective surgery after a single episode of diverticulitis is still a reasonable recommendation (WGO) | C | IIa |
| | Elective sigmoid resection may not be necessary after any specific number of episodes of uncomplicated diverticulitis or with any definite age thresholds (SSAT) | | |
| Immunocompromised Patients | | | |
| Surgical resection | | | |
| Evidence review | Cohort studies indicate that immunocompromised patients have higher risk of complicated recurrence, perforation, and emergency surgery, resulting in a low threshold for operation ^{97,98} ; however, risk of postoperative morbidity and mortality is also higher among these patients ⁹⁴ | C | IIb |
| Practice guidelines ^c | Immunocompromised patients are more likely to present with perforation and to fail medical management, so a lower threshold for urgent and for elective surgery should be applied to them (ASCRS) | C | IIa |
| | Immunocompromised patients (including using steroids and immunosuppressive agents or having diabetes, renal failure, malignancy, cirrhosis) have increased risk of freed perforation, increased need for surgery (WGO) | | |

Abbreviations: ASCRS, American Society of Colon and Rectal Surgeons; SSAT, Society for Surgery of the Alimentary Tract; WGO, World Gastroenterology Organization.

^a Level of recommendations are based on Gibbons et al.⁶ The current practice guidelines are based on Rafferty et al,⁴ SSAT practice guidelines,¹⁰¹ and the WGO practice guidelines.¹⁰²

^b For the definition of the levels of evidence and class, see the Table 1 footnotes.

^c ASCRS represents 2006^{4,101,102}; SSAT, 2007¹⁰¹; and WGO, 2007 practice guidelines.¹⁰²

sequent episodes did not increase the risk of major complications of diverticulitis.⁷⁴ Third, complicated diverticulitis most commonly occurred during the first episode, rather than during recurrent episodes.⁷⁵⁻⁷⁸ Fourth, 5% to 25% of postoperative patients had recurrent or unresolved abdominal symptoms.^{33,79-84}

Patient Characteristics

Although several articles found modestly higher rates of recurrence and need for resection among patients younger than 50 years,^{71,72,85,86} most did not document a greater likelihood of perforation or worse outcomes among this cohort.⁸⁷⁻⁹³ Young patients do have a longer life expectancy (thus, increased potential for future episodes and more to gain from prevention) and lower operative risk.⁹⁴⁻⁹⁶ In general, however, the data supported decision making for operative therapy based on the severity of symptoms and complexity of the disease rather than the age of the patient.^{72,85} Exceptions to the current rule are immunosuppressed patients, such as those with solid organ transplants, collagen vascular diseases, steroid use, malnutrition, and chronic renal failure, who had 5-fold greater risk of complicated recurrence and perforation compared with nonimmunosuppressed patients (36% vs 7%).^{97,98} Thus, the threshold for immunosuppressed patients to proceed with elective resection after 1 or more episodes is lower, provided that the surgical risk due to these same comorbidities is not prohibitive.⁹⁴

Discussion

Sigmoid diverticulitis is an increasingly common and costly disease endemic in industrialized nations. Between 1998 and 2005, US hospital admissions for diverticulitis increased by 26% and elective operations by 29%.⁹⁹ As incidence rates are increasing, the understanding and management of sigmoid diverticulitis is evolving. Collectively, several recent studies indicate a pathogenetic role for inflammation in diverticulitis that may be similar to that of irritable bowel syndrome, inflammatory bowel disease, or both, based on common histologic findings such as granulomas, infiltrating lymphocytes, TNF- α , histamine, and matrix metalloproteinases. However, studies of the etiology and pathophysiology of diverticulitis were limited by a focus on associative rather than causal pathways. Similarly, while lifestyle alterations were often recommended in response to the first incidence of diverticulitis, we found no studies testing the effect of lifestyle modifications on the disease course.

The risk of recurrence among patients with uncomplicated diverticulitis was approximately 13% to 36% and the risk of future emergency surgery was approximately 4% to 7%. Therefore, the available data do not support a routine policy of prophylactic sigmoidectomy on clinical grounds alone (Figure). In addition, given the possibility of shifting morphology,³⁶ a prophylactic resection may

Table 3. Level of Recommendation for Systematic Review of Recent Literature vs Current Practice Guidelines for Management of Chronic Sigmoid Diverticulitis

| Intervention | Current Evidence Review and Guidelines ^a | Recommendation ^b | |
|------------------------------------|--|-----------------------------|-------|
| | | Level | Class |
| Fiber supplementation | | | |
| Evidence review | A systematic review of fiber vs rifaximin + fiber found that the combined therapy was significantly more effective in obtaining symptom relief at 1 y; Use of fiber was not compared with nonuse ⁴⁰ | C | IIb |
| Practice guidelines | Not addressed | | |
| Antibiotic use | | | |
| Evidence review | One study found 90% resolution of symptoms among patients using rifaximin over 1 y; Use of rifaximin alone was not compared with nonuse ⁵² | C | IIb |
| Practice guidelines | Not addressed | | |
| Probiotics | | | |
| Evidence review | A trial of 83 patients randomized to receive oral polybacterial lysate vs placebo reported significantly less abdominal pain, bloating, and fever ⁴⁵ | B | IIa |
| Practice guidelines | Not addressed | | |
| Mesalamine | | | |
| Evidence review | A randomized comparison of mesalamine vs rifaximin demonstrated significantly reduced symptoms after 6-12 mo of high-dose cyclic mesalamine. ⁴³ | B | IIa |
| Practice guidelines | Not addressed | | |
| Confirmation of diagnosis | | | |
| Evidence review | Most cohort studies reported no benefit of colonoscopy among patients following acute diverticulitis ⁶⁵⁻⁶⁸ ; however, several large cohort studies reported a 2%-3% incidence of cancer and up to 26% incidence of adenoma ^{34,63,64} | C | IIa |
| Practice guidelines ^c | Colonoscopy (or contrast enema + sigmoidoscopy) should be uniformly recommended after recovery from acute diverticulitis (ASCRS) | C | I |
| | Care must be taken to exclude other diagnoses (IBS, ischemic colitis) (WGO) | C | IIa |
| | Colonoscopy or barium enema is indicated 6-8 wk following hospital discharge to document the extent of diverticula and exclude cancer (SSAT) | C | I |
| Elective Surgical Resection | | | |
| Evidence review | No study prospectively compared resection to nonresection for chronic diverticulitis. However, several cohort studies examined aspects of chronic disease. Specifically, 5%-25% of postoperative patients had recurrent or unresolved abdominal symptoms ^{33,79-84} | C | IIb |
| Practice guidelines ^c | The decision to recommend surgery should be influenced by whether there are persistent symptoms after the acute episode (ASCRS) | C | I |
| | Sigmoid resection provides complete resolution of smoldering diverticulitis in 70% of cases (WGO) | C | IIa |

Abbreviations: ASCRS, American Society of Colon and Rectal Surgeons; IBS, irritable bowel syndrome; SSAT, Society for Surgery of the Alimentary Tract; WGO, World Gastroenterology Organization.

^a Level of recommendations are based on Gibbons et al.⁶ The current practice guidelines are based on Rafferty et al,⁴ SSAT practice guidelines,¹⁰¹ and the WGO practice guidelines.¹⁰²

^b For the definition of the levels of evidence and class, see the Table 1 footnotes.

^c ASCRS represents 2006^{4,101,102}; SSAT, 2007¹⁰¹; and WGO, 2007 practice guidelines.¹⁰²

actually miss the site of future inflammation, although these data should be confirmed in a larger cohort. Recent data also suggest that combination medical therapy, particularly rifaximin and mesalamine, may contribute to reduced symptoms in chronic disease. Further investigation into commonalities with irritable bowel syndrome and inflammatory bowel disease may provide insight and more opportunities for crossover of medical therapies.

Whether and when to perform elective surgery for chronic or recurrent episodes of uncomplicated diverticulitis remain controversial topics. The traditional recommendation for surgical resection after 2 such episodes was based on outdated evidence suggesting that the success of nonoperative treatment diminished with each subsequent recurrence.¹⁰⁰ Patients were told that an elective operation would permit primary anastomosis, whereas a potential

emergency would necessitate fecal diversion with a colostomy. This recommendation has been challenged by more recent natural history information and recent studies of operative vs nonoperative management.

The current clinical practice guidelines^{4,101,102} were largely written for a surgical audience and may be oriented toward the patient population seen by surgeons (Table 2 and Table 3). However, no corresponding clinical practice guideline specifically targets primary care clinicians who encounter a different spectrum of disease in the primary care setting. Despite current clinical guidelines, we found minimal data to support the prevention of recurrence with fiber ingestion. We found that immunocompromised patients did not have a higher risk of mortality with recurrence. Nor did elective surgery always prevent recurrence or treat chronic disease. Up to 25% of pa-

tients who underwent an operation for chronic diverticulitis had no sustained postoperative symptom relief. Finally, we found mixed data regarding the utility of routine postinflammation colonoscopy. In spite of these data, practice parameters from surgical societies^{4,10,3} stipulate flexible endoscopy to distinguish diverticulitis from other causes of segmental colitis (eg, cancer, Crohn disease, and ischemic colitis).

Our review is subject to a number of limitations which should be noted. Throughout the review, the level of evidence was lower grade, thereby limiting our interpretation and conclusions. Most importantly, studies of diverticulitis were limited by the lack of a standard terminology for aspects of the disease, which resulted in some difficulty commenting across studies. For example, some articles clearly distinguished between complicated and uncomplicated diverticulitis while others did not. Some referred only to diverticular disease and did not discuss diverticulitis per se. Very few studies included outpatient data, and therefore we were unable to draw conclusions about resolution or recurrence of symptoms in this population.

In spite of these limitations, population-based data indicate that peritonitis and sepsis are rare complications of incident diverticulitis and are even more rare complications of recurrent disease. Therefore, although septic peritonitis remains a clear indication for urgent operation, there are few other indications for an urgent colectomy with or without colostomy formation. Moreover, several recommendations meant to reduce the risk of recurrence were based on associations with an index episode and have not been tested longitudinally. Future studies of causality and interventions to reduce recurrence are needed, especially given the increasing incidence of diverticulitis.

Conclusions

In summary, although decision making should be based on individual clinical characteristics and patient preferences, the current data support a substantially reduced role for aggressive antibiotic and surgical intervention in recurrent and chronic diverticulitis.

ARTICLE INFORMATION

Author Contributions: Dr Morris had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Morris, Regenbogen, Hardiman.

Acquisition of data: Morris, Regenbogen, Hardiman.

Analysis and interpretation of data: All authors.

Drafting of the manuscript: All authors.

Critical revision of the manuscript for important intellectual content: All authors.

Administrative, technical, or material support: All authors.

Conflict of Interest Disclosures: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

Submissions: We encourage authors to submit papers for consideration as a Review. Please contact Mary McGrae McDermott, MD, at mdm608@northwestern.edu.

REFERENCES

- Wong WD, Wexner SD, Lowry A, et al; American Society of Colon and Rectal Surgeons. Practice parameters for the treatment of sigmoid diverticulitis—supporting documentation. *Dis Colon Rectum*. 2000;43(3):290-297.
- Ambrosetti P, Becker C, Terrier F. Colonic diverticulitis: impact of imaging on surgical management—a prospective study of 542 patients. *Eur Radiol*. 2002;12(5):1145-1149.
- Tursi A, Joseph RE, Streck P. Expanding applications: the potential usage of 5-aminosalicylic acid in diverticular disease. *Dig Dis Sci*. 2011;56(11):3112-3121.
- Rafferty J, Shellito P, Hyman NH, Buie WD; Standards Committee of American Society of Colon and Rectal Surgeons. Practice parameters for sigmoid diverticulitis. *Dis Colon Rectum*. 2006;49(7):939-944.
- Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting: Meta-analysis of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000;283(15):2008-2012.
- Gibbons RJ, Smith S, Antman E; American College of Cardiology; American Heart Association. American College of Cardiology/American Heart Association clinical practice guidelines, I: where do they come from? *Circulation*. 2003;107(23):2979-2986.
- Guagnini F, Valenti M, Mukenge S, et al. Neural contractions in colonic strips from patients with diverticular disease: role of endocannabinoids and substance P. *Gut*. 2006;55(7):946-953.
- Maselli MA, Piepoli AL, Guerra V, et al. Colonic smooth muscle responses in patients with diverticular disease of the colon: effect of the NK2 receptor antagonist SR48968. *Dig Liver Dis*. 2004;36(5):348-354.
- Golder M, Burleigh DE, Ghali L, et al. Longitudinal muscle shows abnormal relaxation responses to nitric oxide and contains altered levels of NOS1 and elastin in uncomplicated diverticular disease. *Colorectal Dis*. 2007;9(3):218-228.
- Iwase H, Sadahiro S, Mukoyama S, Makuuchi H, Yasuda M. Morphology of myenteric plexuses in the human large intestine: comparison between large intestines with and without colonic diverticula. *J Clin Gastroenterol*. 2005;39(8):674-678.
- Böttner M, Wedel T. Abnormalities of neuromuscular anatomy in diverticular disease. *Dig Dis*. 2012;30(1):19-23.
- Costedio MM, Coates MD, Danielson AB, et al. Serotonin signaling in diverticular disease. *J Gastrointest Surg*. 2008;12(8):1439-1445.
- Morris CR, Harvey IM, Stebbings WS, Speakman CT, Kennedy HJ, Hart AR. Do calcium channel blockers and antimuscarinics protect against perforated colonic diverticular disease? a case control study. *Gut*. 2003;52(12):1734-1737.
- Haboubi NY, Alqudah M. Pathology and pathogenesis of diverticular disease and patterns of colonic mucosal changes overlying the diverticula. *Dig Dis*. 2012;30(1):29-34.
- Goldstein NS, Leon-Armin C, Mani A. Crohn's colitis-like changes in sigmoid diverticulitis specimens is usually an idiosyncratic inflammatory response to the diverticulosis rather than Crohn's colitis. *Am J Surg Pathol*. 2000;24(5):668-675.
- Tursi A, Elisei W, Brandimarte G, et al. Mucosal expression of basic fibroblastic growth factor, Syndecan 1 and tumor necrosis factor-alpha in diverticular disease of the colon: a case-control study. *Neurogastroenterol Motil*. 2012;24(9):836-e396.
- von Rahden BH, Jurowich C, Kircher S, et al. Allergic predisposition, histamine and histamine receptor expression (H1R, H2R) are associated with complicated courses of sigmoid diverticulitis. *J Gastrointest Surg*. 2012;16(1):173-182, discussion 182.
- Altadill A, Eiró N, González LO, et al. Comparative analysis of the expression of metalloproteases and their inhibitors in resected Crohn's disease and complicated diverticular disease. *Inflamm Bowel Dis*. 2012;18(1):120-130.
- Crowe FL, Appleby PN, Allen NE, Key TJ. Diet and risk of diverticular disease in Oxford cohort of European Prospective Investigation into Cancer and Nutrition (EPIC): prospective study of British vegetarians and non-vegetarians. *BMJ*. 2011;343:d4131.
- Peery AF, Barrett PR, Park D, et al. A high-fiber diet does not protect against asymptomatic diverticulosis. *Gastroenterology*. 2012;142(2):266-272.
- Strate LL, Liu YL, Syngal S, Aldoori WH, Giovannucci EL. Nut, corn, and popcorn consumption and the incidence of diverticular disease. *JAMA*. 2008;300(8):907-914.
- Dobbins C, Defontgalland D, Duthie G, Wattchow DA. The relationship of obesity to the complications of diverticular disease. *Colorectal Dis*. 2006;8(1):37-40.
- Hjern F, Wolk A, Håkansson N. Obesity, physical inactivity, and colonic diverticular disease requiring hospitalization in women: a prospective cohort study. *Am J Gastroenterol*. 2012;107(2):296-302.

24. Rosemar A, Angerås U, Rosengren A. Body mass index and diverticular disease: a 28-year follow-up study in men. *Dis Colon Rectum*. 2008;51(4):450-455.
25. Strate LL, Liu YL, Aldoori WH, Syngal S, Giovannucci EL. Obesity increases the risks of diverticulitis and diverticular bleeding. *Gastroenterology*. 2009;136(1):115-122.
26. Hjern F, Wolk A, Håkansson N. Smoking and the risk of diverticular disease in women. *Br J Surg*. 2011;98(7):997-1002.
27. Strate LL, Liu YL, Aldoori WH, Giovannucci EL. Physical activity decreases diverticular complications. *Am J Gastroenterol*. 2009;104(5):1221-1230.
28. Williams PT. Incident diverticular disease is inversely related to vigorous physical activity. *Med Sci Sports Exerc*. 2009;41(5):1042-1047.
29. Morris CR, Harvey IM, Stebbings WS, Speakman CT, Kennedy HJ, Hart AR. Anti-inflammatory drugs, analgesics and the risk of perforated colonic diverticular disease. *Br J Surg*. 2003;90(10):1267-1272.
30. Strate LL, Liu YL, Huang ES, Giovannucci EL, Chan AT. Use of aspirin or nonsteroidal anti-inflammatory drugs increases risk for diverticulitis and diverticular bleeding. *Gastroenterology*. 2011;140(5):1427-1433.
31. von Rahden BH, Kircher S, Thiery S, et al. Association of steroid use with complicated sigmoid diverticulitis: potential role of activated CD68+/CD163+ macrophages. *Langenbecks Arch Surg*. 2011;396(6):759-768.
32. Goh H, Bourne R. Non-steroidal anti-inflammatory drugs and perforated diverticular disease: a case-control study. *Ann R Coll Surg Engl*. 2002;84(2):93-96.
33. Binda GA, Arezzo A, Serventi A, et al; Italian Study Group on Complicated Diverticulosis (GISDIC). Multicentre observational study of the natural history of left-sided acute diverticulitis [published correction appears in *Br J Surg*. 2012;99(4):600]. *Br J Surg*. 2012;99(2):276-285.
34. Broderick-Villa G, Burchette RJ, Collins JC, Abbas MA, Haigh PI. Hospitalization for acute diverticulitis does not mandate routine elective colectomy. *Arch Surg*. 2005;140(6):576-581.
35. Humes DJ, West J. Role of acute diverticulitis in the development of complicated colonic diverticular disease and 1-year mortality after diagnosis in the UK: population-based cohort study. *Gut*. 2012;61(1):95-100.
36. Gervaz P, Platon A, Widmer L, Ambrosetti P, Poletti PA. A clinical and radiological comparison of sigmoid diverticulitis episodes 1 and 2. *Colorectal Dis*. 2012;14(4):463-468.
37. Nelson RS, Ewing BM, Wengert TJ, Thorson AG. Clinical outcomes of complicated diverticulitis managed nonoperatively. *Am J Surg*. 2008;196(6):969-972.
38. Humes DJ, Simpson J, Neal KR, Scholefield JH, Spiller RC. Psychological and colonic factors in painful diverticulosis. *Br J Surg*. 2008;95(2):195-198.
39. Alonso S, Pera M, Parés D, et al. Outpatient treatment of patients with uncomplicated acute diverticulitis. *Colorectal Dis*. 2010;12(10 online):e278-e282.
40. Bianchi M, Festa V, Moretti A, et al. Long-term therapy with rifaximin in the management of uncomplicated diverticular disease. *Aliment Pharmacol Ther*. 2011;33(8):902-910.
41. Brandimarte G, Tursi A. Rifaximin plus mesalazine followed by mesalazine alone is highly effective in obtaining remission of symptomatic uncomplicated diverticular disease. *Med Sci Monit*. 2004;10(5):P170-P173.
42. Chabok A, Pählman L, Hjern F, Haapaniemi S, Smedh K; AVOD Study Group. Randomized clinical trial of antibiotics in acute uncomplicated diverticulitis. *Br J Surg*. 2012;99(4):532-539.
43. Comparato G, Fanigliuolo L, Cavallaro LG, et al. Prevention of complications and symptomatic recurrences in diverticular disease with mesalazine: a 12-month follow-up. *Dig Dis Sci*. 2007;52(11):2934-2941.
44. de Korte N, Unlü C, Boermeester MA, Cuesta MA, Vrouenraets BC, Stockmann HB. Use of antibiotics in uncomplicated diverticulitis. *Br J Surg*. 2011;98(6):761-767.
45. Dughera L, Serra AM, Battaglia E, Tibaudi D, Navino M, Emanuelli G. Acute recurrent diverticulitis is prevented by oral administration of a polybacterial lysate suspension. *Minerva Gastroenterol Dietol*. 2004;50(2):149-153.
46. Hjern F, Josephson T, Altman D, et al. Conservative treatment of acute colonic diverticulitis: are antibiotics always mandatory? *Scand J Gastroenterol*. 2007;42(1):41-47.
47. Mizuki A, Nagata H, Tatemichi M, et al. The out-patient management of patients with acute mild-to-moderate colonic diverticulitis. *Aliment Pharmacol Ther*. 2005;21(7):889-897.
48. Moya P, Arroyo A, Pérez-Legaz J, et al. Applicability, safety and efficiency of outpatient treatment in uncomplicated diverticulitis. *Tech Coloproctol*. 2012;16(4):301-307.
49. Ribas Y, Bombardó J, Aguilar F, et al. Prospective randomized clinical trial assessing the efficacy of a short course of intravenously administered amoxicillin plus clavulanic acid followed by oral antibiotic in patients with uncomplicated acute diverticulitis. *Int J Colorectal Dis*. 2010;25(11):1363-1370.
50. Ridgway PF, Latif A, Shabbir J, et al. Randomized controlled trial of oral vs intravenous therapy for the clinically diagnosed acute uncomplicated diverticulitis. *Colorectal Dis*. 2009;11(9):941-946.
51. Shabanzadeh DM, Wille-Jørgensen P. Antibiotics for uncomplicated diverticulitis. *Cochrane Database Syst Rev*. 2012;11:CD009092.
52. Tursi A, Brandimarte G, Daffinà R. Long-term treatment with mesalazine and rifaximin versus rifaximin alone for patients with recurrent attacks of acute diverticulitis of colon. *Dig Liver Dis*. 2002;34(7):510-515.
53. Ünlü C, Daniels L, Vrouenraets BC, Boermeester MA. A systematic review of high-fibre dietary therapy in diverticular disease. *Int J Colorectal Dis*. 2012;27(4):419-427.
54. Schug-Pass C, Geers P, Hügel O, Lippert H, Köckerling F. Prospective randomized trial comparing short-term antibiotic therapy versus standard therapy for acute uncomplicated sigmoid diverticulitis. *Int J Colorectal Dis*. 2010;25(6):751-759.
55. Regenbogen SE, Hardiman KM, Hendren S, Morris AM. Surgery for diverticulitis in the 21st century: a systematic review. *JAMASurg*. 10.1001/jamasurg.2013.5477.
56. Costi R, Cauchy F, Le Bian A, Honart JF, Creuze N, Smadja C. Challenging a classic myth: pneumoperitoneum associated with acute diverticulitis is not an indication for open or laparoscopic emergency surgery in hemodynamically stable patients: a 10-year experience with a nonoperative treatment. *Surg Endosc*. 2012;26(7):2061-2071.
57. Dharmarajan S, Hunt SR, Birnbaum EH, Fleshman JW, Mutch MG. The efficacy of nonoperative management of acute complicated diverticulitis. *Dis Colon Rectum*. 2011;54(6):663-671.
58. Ambrosetti P. Value of CT for acute left-colonic diverticulitis: the surgeon's view. *Dig Dis*. 2012;30(1):51-55.
59. Ambrosetti P, Chautems R, Soravia C, Peiris-Waser N, Terrier F. Long-term outcome of mesocolic and pelvic diverticular abscesses of the left colon: a prospective study of 73 cases. *Dis Colon Rectum*. 2005;48(4):787-791.
60. Kaiser AM, Jiang JK, Lake JP, et al. The management of complicated diverticulitis and the role of computed tomography. *Am J Gastroenterol*. 2005;100(4):910-917.
61. Poletti PA, Platon A, Rutschmann O, et al. Acute left colonic diverticulitis: can CT findings be used to predict recurrence? *AJR Am J Roentgenol*. 2004;182(5):1159-1165.
62. Shaikh S, Krukowski ZH. Outcome of a conservative policy for managing acute sigmoid diverticulitis. *Br J Surg*. 2007;94(7):876-879.
63. Lahat A, Yanai H, Sakhnini E, Menachem Y, Bar-Meir S. Role of colonoscopy in patients with persistent acute diverticulitis. *World J Gastroenterol*. 2008;14(17):2763-2766.
64. Lau KC, Spilsbury K, Farooque Y, et al. Is colonoscopy still mandatory after a CT diagnosis of left-sided diverticulitis: can colorectal cancer be confidently excluded? *Dis Colon Rectum*. 2011;54(10):1265-1270.
65. Bar-Meir S, Lahat A, Melzer E. Role of endoscopy in patients with diverticular disease. *Dig Dis*. 2012;30(1):60-63.
66. Lam TJ, Meurs-Szojda MM, Gundlach L, et al. There is no increased risk for colorectal cancer and adenomas in patients with diverticulitis: a retrospective longitudinal study. *Colorectal Dis*. 2010;12(11):1122-1126.
67. Schmilovitz-Weiss H, Yalunin E, Boaz M, et al. Does a colonoscopy after acute diverticulitis affect its management? a single center experience. *J Clin Gastroenterol*. 2012;46(4):317-320.
68. Westwood DA, Eglinton TW, Frizelle FA. Routine colonoscopy following acute uncomplicated diverticulitis. *Br J Surg*. 2011;98(11):1630-1634.
69. Eglinton T, Nguyen T, Raniga S, Dixon L, Dobbs B, Frizelle FA. Patterns of recurrence in patients with acute diverticulitis. *Br J Surg*. 2010;97(6):952-957.
70. Hall JF, Roberts PL, Ricciardi R, et al. Long-term follow-up after an initial episode of diverticulitis:

what are the predictors of recurrence? *Dis Colon Rectum*. 2011;54(3):283-288.

71. Anaya DA, Flum DR. Risk of emergency colectomy and colostomy in patients with diverticular disease. *Arch Surg*. 2005;140(7):681-685.
72. Guzzo J, Hyman N. Diverticulitis in young patients: is resection after a single attack always warranted? *Dis Colon Rectum*. 2004;47(7):1187-1190.
73. Janes S, Meagher A, Frizelle FA. Elective surgery after acute diverticulitis. *Br J Surg*. 2005;92(2):133-142.
74. Chapman JR, Dozois EJ, Wolff BG, Gullerud RE, Larson DR. Diverticulitis: a progressive disease? do multiple recurrences predict less favorable outcomes? *Ann Surg*. 2006;243(6):876-830.
75. Issa N, Dreznik Z, Dueck DS, et al. Emergency surgery for complicated acute diverticulitis. *Colorectal Dis*. 2009;11(2):198-202.
76. Pittet O, Kotzampassakis N, Schmidt S, Denys A, Demartines N, Calmes JM. Recurrent left colonic diverticulitis episodes: more severe than the initial diverticulitis? *World J Surg*. 2009;33(3):547-552.
77. Ritz JP, Lehmann KS, Frericks B, Stroux A, Buhr HJ, Holmer C. Outcome of patients with acute sigmoid diverticulitis: multivariate analysis of risk factors for free perforation. *Surgery*. 2011;149(5):606-613.
78. Chapman J, Davies M, Wolff B, et al. Complicated diverticulitis: is it time to rethink the rules? *Ann Surg*. 2005;242(4):576-581.
79. Andeweg C, Peters J, Bleichrodt R, van Goor H. Incidence and risk factors of recurrence after surgery for pathology-proven diverticular disease. *World J Surg*. 2008;32(7):1501-1506.
80. Ambrosetti P, Francis K, Weintraub D, Weintraub J. Functional results following elective laparoscopic sigmoidectomy after CT-proven diagnosis of acute diverticulitis evaluation of 43 patients and review of the literature. *J Gastrointest Surg*. 2007;11(6):767-772.
81. Egger B, Peter MK, Candinas D. Persistent symptoms after elective sigmoid resection for diverticulitis. *Dis Colon Rectum*. 2008;51(7):1044-1048.
82. Forgione A, Leroy J, Cahill RA, et al. Prospective evaluation of functional outcome after laparoscopic sigmoid colectomy. *Ann Surg*. 2009;249(2):218-224.
83. Käser SA, Glauser PM, Basilicata G, Müller DA, Maurer CA. Timing of resectosigmoid resection for diverticular disease: the patient's view. *Colorectal Dis*. 2012;14(3):e111-e116.
84. Pasternak I, Wiedemann N, Basilicata G, Melcher GA. Gastrointestinal quality of life after laparoscopic-assisted sigmoidectomy for diverticular disease. *Int J Colorectal Dis*. 2012;27(6):781-787.
85. Faria GR, Almeida AB, Moreira H, Pinto-de-Sousa J, Correia-da-Silva P, Pimenta AP. Acute diverticulitis in younger patients: any rationale for a different approach? *World J Gastroenterol*. 2011;17(2):207-212.
86. Lahat A, Menachem Y, Avidan B, et al. Diverticulitis in the young patient—is it different? *World J Gastroenterol*. 2006;12(18):2932-2935.
87. Biondo S, Parés D, Martí Ragué J, Kreisler E, Fracalvieri D, Jaurrieta E. Acute colonic diverticulitis in patients under 50 years of age. *Br J Surg*. 2002;89(9):1137-1141.
88. Hjern F, Josephson T, Altman D, Holmström B, Johansson C. Outcome of younger patients with acute diverticulitis. *Br J Surg*. 2008;95(6):758-764.
89. Kotzampassakis N, Pittet O, Schmidt S, Denys A, Demartines N, Calmes JM. Presentation and treatment outcome of diverticulitis in younger adults: a different disease than in older patients? *Dis Colon Rectum*. 2010;53(3):333-338.
90. Lopez-Borao J, Kreisler E, Millan M, et al. Impact of age on recurrence and severity of left colonic diverticulitis. *Colorectal Dis*. 2012;14(7):e407-e412.
91. Mäkelä JT, Kiviniemi HO, Laitinen ST. Acute sigmoid diverticulitis in young patients. *Hepatogastroenterology*. 2009;56(94-95):1382-1387.
92. Nelson RS, Velasco A, Mukesh BN. Management of diverticulitis in younger patients. *Dis Colon Rectum*. 2006;49(9):1341-1345.
93. Ritz JP, Lehmann KS, Stroux A, Buhr HJ, Holmer C. Sigmoid diverticulitis in young patients—a more aggressive disease than in older patients? *J Gastrointest Surg*. 2011;15(4):667-674.
94. Pessaix P, Muscari F, Ouellet JF, et al. Risk factors for mortality and morbidity after elective sigmoid resection for diverticulitis: prospective multicenter multivariate analysis of 582 patients. *World J Surg*. 2004;28(1):92-96.
95. Oomen JL, Engel AF, Cuesta MA. Mortality after acute surgery for complications of diverticular disease of the sigmoid colon is almost exclusively due to patient related factors. *Colorectal Dis*. 2006;8(2):112-119.
96. Lidsky ME, Thacker JK, Lagoo-Deenadayalan SA, Scarborough JE. Advanced age is an independent predictor for increased morbidity and mortality after emergent surgery for diverticulitis. *Surgery*. 2012;152(3):465-472.
97. Klarenbeek BR, Samuels M, van der Wal MA, van der Peet DL, Meijerink WJ, Cuesta MA. Indications for elective sigmoid resection in diverticular disease. *Ann Surg*. 2010;251(4):670-674.
98. Yoo PS, Garg R, Salamone LF, Floch MH, Rosenthal R, Longo WE. Medical comorbidities predict the need for colectomy for complicated and recurrent diverticulitis. *Am J Surg*. 2008;196(5):710-714.
99. Etzioni DA, Mack TM, Beart RW Jr, Kaiser AM. Diverticulitis in the United States: 1998-2005: changing patterns of disease and treatment. *Ann Surg*. 2009;249(2):210-217.
100. Parks TG. Natural history of diverticular disease of the colon: a review of 521 cases. *BMJ*. 1969;4(5684):639-642.
101. Society for Surgery of the Alimentary Tract. Patient care guidelines. <http://www.ssat.com/cgi-bin/divert.cgi>. Board approved October 8, 2007. Accessed April 4, 2013.
102. World Gastroenterology Organisation. Practice guidelines—diverticular disease. <http://www.worldgastroenterology.org/diverticular-disease.html>. Accessed April 4, 2013.
103. Fozard JB, Armitage NC, Schofield JB, Jones OM; Association of Coloproctology of Great Britain and Ireland. ACPGBI position statement on elective resection for diverticulitis. *Colorectal Dis*. 2011;13(suppl 3):1-11.