

## Gastroesophageal Reflux Disease

Heartburn, acid reflux, and acid indigestion are all terms used to describe a burning feeling in the chest or a backwash of stomach contents into the mouth or the esophagus (the tube that connects the throat to the stomach).

If you have these symptoms more than twice a week for more than a few weeks, you may be diagnosed as having **gastroesophageal reflux disease (GERD)**.

### Symptoms

The most common symptoms of GERD are a burning feeling in the chest (heartburn) and regurgitating food or liquid into the throat. Other symptoms include dry, chronic cough; wheezing or asthma; sore throat or hoarseness; and tooth erosion.

*If you have pain or pressure in your chest, shortness of breath, nausea or vomiting, or back or arm pain, you might be experiencing a heart attack. Seek emergency medical attention right away.*

### Diagnosis

Your doctor will most likely diagnose GERD based on your symptoms and whether your symptoms improve with medication. Other tests that are sometimes used include

- Endoscopy, in which the doctor uses a video camera to look at the esophagus
- X-ray scans of the upper gastrointestinal tract to look for narrowing of the esophagus or other problems
- Manometry to measure the muscle contractions of the esophagus

### Treatment

There are a number of treatments you can try at home. You may need to use 1 or more treatments on an ongoing basis.


- **Medication:** Over-the-counter medications include antacids and 2 types of medicines that reduce stomach acid: H<sub>2</sub> receptor blockers (such as ranitidine) and proton pump inhibitors (such as omeprazole).
- **Changes in diet:** Some people find it helpful to avoid eating or drinking chocolate, coffee, peppermint, greasy or spicy foods, tomato products, and alcohol.

Other treatments include quitting smoking, avoiding overeating, avoiding eating within 2 to 3 hours before bedtime, and trying to lose weight if you are overweight. If your symptoms are worse at night, it may help to prop up the head of your bed about 6 to 8 inches. Using extra pillows will not help GERD.

### Surgery

Sometimes doctors recommend surgery for GERD, most commonly laparoscopic fundoplication. This procedure involves folding the top

Gastroesophageal reflux disease



Reflux occurs when your **esophageal sphincter** does not close properly. This allows food or acid **reflux** to wash back up into your **esophagus**, causing irritation. Treatments include over-the-counter and prescription antacid medications, diet and lifestyle changes, and surgery.

portion of the stomach over the opening of the esophagus to prevent backwash of contents. In some cases, the doctor may recommend weight loss surgery.

### Complications

Left untreated, GERD can cause problems in the esophagus, including ulcers, swallowing problems, or Barrett esophagus, a condition that increases risk of esophageal cancer.

#### FOR MORE INFORMATION

- National Library of Medicine  
[digestive.niddk.nih.gov/ddiseases/pubs/gerd](http://digestive.niddk.nih.gov/ddiseases/pubs/gerd)

**+** To find this and previous JAMA Patient Pages, go to the Patient Page link on JAMA's website at [jama.com](http://jama.com). Many are available in English and Spanish. An article on laparoscopic antireflux surgery vs esomeprazole treatment for chronic GERD was published in the May 18, 2011, issue of JAMA. A clinical review of Barrett esophagus and risk of esophageal cancer and accompanying online video were published in the August 14, 2013, issue of JAMA.

**Author:** Deborah Tolmach Sugerman, MSW

**Conflict of Interest Disclosures:** The author has completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none were reported.

**Source:** Katz PO, Gerson LB, Vela MF. Guidelines for the diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol*. 2013;108:308-328.

The JAMA Patient Page is a public service of JAMA. The information and recommendations appearing on this page are appropriate in most instances, but they are not a substitute for medical diagnosis. For specific information concerning your personal medical condition, JAMA suggests that you consult your physician. This page may be photocopied noncommercially by physicians and other health care professionals to share with patients. To purchase bulk reprints, call 312/464-0776.